



Understanding Side Effects

This brochure will help you understand and talk about your side effects with your healthcare team.

REVLIMID is only available through a restricted distribution program called Lenalidomide REMS.

Please see REVLIMID Important Safety Information throughout and on pages 2-15, as well as full [Prescribing Information](#), including Boxed WARNINGS regarding risk to unborn babies, risk of low blood counts, and blood clots, and [Medication Guide](#).



“I try to listen to my own body and communicate with my healthcare team about how I’m feeling.”

– Julie, patient with MM

WHY IS IT IMPORTANT TO LEARN ABOUT SIDE EFFECTS?

While taking REVLIMID® (lenalidomide), you may experience side effects. This guide provides tips and questions for you to discuss with your doctor in the event a side effect occurs.

Always talk with your healthcare team about any side effects you experience.

Important Safety Information

What is REVLIMID® (lenalidomide)?

REVLIMID is a prescription medicine used to treat adults with multiple myeloma (MM) in combination with the medicine dexamethasone, or as maintenance treatment after autologous hematopoietic stem cell transplantation (a type of stem cell transplant that uses your own stem cells).

Please see Important Safety Information throughout and full [Prescribing Information](#), including Boxed WARNINGS and [Medication Guide](#).



Why should I talk to my doctor about side effects?

It's important to tell your healthcare team about the side effects you experience in your treatment journey.

Staying on REVLIMID® (lenalidomide) until your disease progresses, or until you experience severe side effects that you cannot tolerate, is also important. Your doctor may adjust your dose or tell you to stop taking REVLIMID if you experience serious side effects during treatment.

How long will I stay on REVLIMID?

Everyone's experience with REVLIMID is different, so it's hard to say exactly how long you'll be on treatment. But it's important to stay on treatment, as patients who continuously took REVLIMID lived longer without their disease progressing.



Make sure you get help right away if any side effects become unbearable.

Important Safety Information

What is REVLIMID® (lenalidomide)? (continued)

REVLIMID should not be used to treat people who have chronic lymphocytic leukemia (CLL) unless they are participants in a controlled clinical trial.

It is not known if REVLIMID is safe and effective in children.

WARNING: Risk to unborn babies, risk of low blood counts and blood clots.

Please see Important Safety Information throughout and full [Prescribing Information](#), including Boxed WARNINGS and [Medication Guide](#).



What are the possible side effects?

SERIOUS SIDE EFFECTS OF REVLIMID® (lenalidomide) INCLUDE:

- Birth defects
- Increased risk of death in people with chronic lymphocytic leukemia (CLL)
- Risk of new cancers (malignancies)
- Severe liver problems (including liver failure and death)
- Severe skin reactions and severe allergic reactions
- Tumor lysis syndrome (TLS)
- Worsening of your tumor (tumor flare reaction)
- Thyroid problems
- Risk of early death in mantle cell lymphoma (MCL)

The full details about these serious side effects can be found under “What are the possible side effects of REVLIMID?” on pages 13-15 of the Important Safety Information. REVLIMID is only available through Lenalidomide REMS.

Important Safety Information (continued)

What is the most important information I should know about REVLIMID?

Before you begin taking REVLIMID, you must read and agree to all of the instructions in the Lenalidomide REMS program. Before prescribing REVLIMID, your healthcare provider will explain the Lenalidomide REMS program to you and have you sign the Patient-Physician Agreement Form.

REVLIMID may cause serious side effects, including:

- **Possible birth defects (deformed babies) or death of an unborn baby.** Females who are pregnant or who plan to become pregnant must not take REVLIMID.

REVLIMID is similar to the medicine thalidomide which is known to cause severe life-threatening birth defects. REVLIMID has not been tested in pregnant females. REVLIMID has harmed unborn animals in animal testing.

Please see Important Safety Information throughout and full [Prescribing Information](#), including Boxed WARNINGS and [Medication Guide](#).



THE MOST COMMON SIDE EFFECTS OF REVLIMID® (lenalidomide) INCLUDE:

- Diarrhea
- Rash
- Nausea
- Constipation
- Tiredness or weakness
- Fever
- Itching
- Swelling of your arms, hands, legs, feet, and skin
- Sleep problems (insomnia)
- Headache
- Muscle cramps or spasms
- Shortness of breath
- Cough, sore throat, and other symptoms of a cold
- Upper respiratory tract infection or bronchitis
- Inflammation of your stomach and intestine (stomach flu)
- Nose bleed
- Shaking or trembling (tremor)
- Joint aches
- Pain in your back or stomach area (abdomen)

These are not all of the possible side effects of REVLIMID.

Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088.

Important Safety Information (continued)

What is the most important information I should know about REVLIMID? (continued)

Females must not get pregnant:

- For at least 4 weeks before starting REVLIMID
- While taking REVLIMID
- During any breaks (interruptions) in your treatment with REVLIMID
- For at least 4 weeks after stopping REVLIMID

Please see Important Safety Information throughout and full [Prescribing Information](#), including Boxed WARNINGS and [Medication Guide](#).



What should I do if I experience side effects?

To help you talk to your doctor about side effects, here are some questions you can ask and tips that may help.

Diarrhea

ASK YOUR DOCTOR:

- How much liquid should I drink each day?
- Is there a medicine I can take to help with the diarrhea?



Drink clear liquids like water or broth.
Avoid caffeine and alcohol.

Important Safety Information (continued)

What is the most important information I should know about REVLIMID® (lenalidomide)? (continued)

Females who can become pregnant:

- Must have pregnancy tests weekly for 4 weeks, then every 4 weeks if your menstrual cycle is regular, or every 2 weeks if your menstrual cycle is irregular.
- If you miss your period or have unusual bleeding, you will need to have a pregnancy test and receive counseling.
- Must agree to use 2 different forms of effective birth control at the same time, for at least 4 weeks before, while taking, during any breaks (interruptions) in your treatment, and for at least 4 weeks after stopping REVLIMID.
- Talk with your healthcare provider to find out about options for effective forms of birth control that you may use to prevent pregnancy before, during, and after treatment with REVLIMID.
- If you had unprotected sex or if you think your birth control has failed, stop taking REVLIMID immediately and call your healthcare provider right away.

Please see Important Safety Information throughout and full [Prescribing Information](#), including Boxed WARNINGS and [Medication Guide](#).



Itching and rash

Some patients have had serious skin reactions that required immediate medical attention or were fatal. If you get any kind of rash, tell your healthcare provider immediately.

ASK YOUR DOCTOR:

- Should I take hot or cold baths?
- Should I avoid the sun?
- Are there any medicines that I can take to help relieve the itching?

Nausea

ASK YOUR DOCTOR:

- Are there foods I should avoid when I have nausea?
- Are there medicines that may help?

Tip

Drink clear liquids. Make sure they're cold and sipped slowly.



REVLIMID® (lenalidomide) is available in different doses, so you may be able to work with your doctor to adjust your dose, depending on side effects.

Important Safety Information (continued)

What is the most important information I should know about REVLIMID? (continued)

If you become pregnant while taking REVLIMID, stop taking it right away and call your healthcare provider. If your healthcare provider is not available, you can call Celgene Customer Care Center at 1-888-423-5436. Healthcare providers and patients should report all cases of pregnancy to:

- FDA MedWatch at 1-800-FDA-1088, and
- Celgene Corporation at 1-888-423-5436.

Please see Important Safety Information throughout and full [Prescribing Information](#), including Boxed WARNINGS and [Medication Guide](#).



Constipation

ASK YOUR DOCTOR:

- Should I drink more water or liquids?
- Should I keep track of my bowel movements?
- Are there any medicines that may help?

Tip

Work with your doctor to see if an exercise plan is appropriate.

Important Safety Information (continued)

What is the most important information I should know about REVLIMID® (lenalidomide)? (continued)

There is a pregnancy exposure registry that monitors the outcomes of females who take REVLIMID during pregnancy, or if their male partner takes REVLIMID and they are exposed during pregnancy. You can enroll in this registry by calling Celgene Corporation at the phone number listed above.

REVLIMID can pass into human semen:

- Males, including those who have had a vasectomy, must always use a latex or synthetic condom during any sexual contact with a pregnant female or a female that can become pregnant while taking REVLIMID, during any breaks (interruptions) in your treatment with REVLIMID, and for up to 4 weeks after stopping REVLIMID.
- Do not have unprotected sexual contact with a female who is or could become pregnant. Tell your healthcare provider if you do have unprotected sexual contact with a female who is or could become pregnant.
- Do not donate sperm while taking REVLIMID, during any breaks (interruptions) in your treatment, and for 4 weeks after stopping REVLIMID. If a female becomes pregnant with your sperm, the baby may be exposed to REVLIMID and may be born with birth defects.

Men: If a female becomes pregnant with your sperm, you should call your HCP right away.

Please see Important Safety Information throughout and full [Prescribing Information](#), including Boxed WARNINGS and [Medication Guide](#).



Fatigue

ASK YOUR DOCTOR:

- Would exercise help me feel less tired?
- Are there medicines that may help?
- If I am already taking a medicine that makes me feel tired, will REVLIMID® (lenalidomide) make me more tired?



It's important to always talk with your healthcare team about your side effects.

Important Safety Information (continued)

What is the most important information I should know about REVLIMID? (continued)

- **Low white blood cells (neutropenia) and low platelets (thrombocytopenia).** REVLIMID causes low white blood cells and low platelets in most people. You may need a blood transfusion or certain medicines if your blood counts drop too low. Your healthcare provider should check your blood counts often, especially during the first several months of treatment with REVLIMID, and then at least monthly. Tell your healthcare provider if you develop any bleeding or bruising during treatment with REVLIMID.

Please see Important Safety Information throughout and full [Prescribing Information](#), including Boxed WARNINGS and [Medication Guide](#).



Fever

ASK YOUR DOCTOR:

- Are there any dietary restrictions I should follow?
- Are there medicines that may help?

Tip

Drink fluids like water, broth, or juice.

Swelling of the limbs and skin

ASK YOUR DOCTOR:

- Should I drink more or less water or liquids?
- Are there medicines that may help?

Cough

ASK YOUR DOCTOR:

- Should I take cough drops?
- Are there medicines that may help?

Tip

Drink fluids.

Please see Important Safety Information throughout and full [Prescribing Information](#), including Boxed WARNINGS and [Medication Guide](#).


Revlimid[®]
(lenalidomide) capsules
2.5 · 5 · 10 · 15 · 20 · 25 mg

Important Safety Information (continued)

What is the most important information I should know about REVLIMID® (lenalidomide)? (continued)

- **Blood clots.** Blood clots in the arteries, veins, and lungs happen more often in people who take REVLIMID. This risk is even higher for people with multiple myeloma who take the medicine dexamethasone with REVLIMID. Heart attacks and strokes also happen more often in people who take REVLIMID with dexamethasone. To reduce this increased risk, most people who take REVLIMID will also take a blood thinner medicine.

Before taking REVLIMID, tell your healthcare provider:

- if you have had a blood clot in the past;
- if you have high blood pressure, smoke, or if you have been told you have a high level of fat in your blood (hyperlipidemia); and
- about all the medicines you take. Certain other medicines can also increase your risk for blood clots

Call your healthcare provider or get medical help right away if you get any of the following during treatment with REVLIMID:

- **Signs or symptoms of a blood clot in the lung, arm, or leg may include:** shortness of breath, chest pain, or arm or leg swelling
- **Signs or symptoms of a heart attack may include:** chest pain that may spread to the arms, neck, jaw, back, or stomach area (abdomen), feeling sweaty, shortness of breath, feeling sick or vomiting
- **Signs or symptoms of stroke may include:** sudden numbness or weakness, especially on one side of the body, severe headache or confusion, or problems with vision, speech, or balance

Who should not take REVLIMID?

Do not take REVLIMID if you:

- **are pregnant, plan to become pregnant, or become pregnant during treatment with REVLIMID. See “What is the most important information I should know about REVLIMID?”**
- are allergic to lenalidomide or any of the ingredients in REVLIMID. See the [Medication Guide](#) for a complete list of ingredients in REVLIMID.

Please see Important Safety Information throughout and full [Prescribing Information](#), including Boxed WARNINGS and [Medication Guide](#).



Important Safety Information (continued)

What should I tell my healthcare provider before taking REVLIMID® (lenalidomide)?

Before you take REVLIMID, tell your healthcare provider about all of your medical conditions, including if you:

- have liver problems
- have kidney problems or receive kidney dialysis treatment
- have thyroid problems
- have had a serious skin rash with thalidomide treatment. You should not take REVLIMID.
- are lactose intolerant. REVLIMID contains lactose.
- are breastfeeding. Do not breastfeed during treatment with REVLIMID. It is not known if REVLIMID passes into your breast milk and can harm your baby.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. REVLIMID and other medicines may affect each other, causing serious side effects. Talk with your healthcare provider before taking any new medicines. Know the medicines you take. Keep a list of them to show your healthcare provider and pharmacist.

How should I take REVLIMID?

Take REVLIMID exactly as prescribed and follow all the instructions of the Lenalidomide REMS program

- Swallow REVLIMID capsules whole, with water, 1 time a day. **Do not open, break, or chew your capsules.**
- **REVLIMID may be taken with or without food.**
- Take REVLIMID at about the same time each day.
- Do not open the REVLIMID capsules or handle them any more than needed. If powder from the REVLIMID capsule comes in contact with:
 - your skin, wash the skin right away with soap and water.
 - inside of your eyes, nose, or mouth, flush well with water.
- If you miss a dose of REVLIMID and it has been less than 12 hours since your regular time, take it as soon as you remember. If it has been more than 12 hours, just skip your missed dose. Do **not** take 2 doses at the same time.

Please see Important Safety Information throughout and full [Prescribing Information](#), including Boxed WARNINGS and [Medication Guide](#).



Important Safety Information (continued)

What should I avoid while taking REVLIMID® (lenalidomide)?

- If you take too much REVLIMID, call your healthcare provider right away.
- See “What is the most important information I should know about REVLIMID?”
- **Females: Do not get pregnant and do not breastfeed while taking REVLIMID.**
- **Males: Do not donate sperm.**
- **Do not share REVLIMID with other people. It may cause birth defects and other serious problems.**
- **Do not donate blood** while you take REVLIMID, during any breaks (interruptions) in your treatment, and for 4 weeks after stopping REVLIMID. If someone who is pregnant gets your donated blood, her baby may be exposed to REVLIMID and may be born with birth defects.

What are the possible side effects of REVLIMID?

REVLIMID can cause serious side effects, including:

- **See “What is the most important information I should know about REVLIMID?”**
- **Increased risk of death in people who have chronic lymphocytic leukemia (CLL).** People with CLL who take REVLIMID have an increased risk of death compared with people who take the medicine chlorambucil. REVLIMID may cause you to have serious heart problems that can lead to death, including atrial fibrillation, heart attack, or heart failure. You should not take REVLIMID if you have CLL unless you are participating in a controlled clinical trial.
- **Risk of new cancers (malignancies).** An increase in new (second) cancers has happened in patients who received REVLIMID and melphalan, or a blood stem cell transplant, including certain blood cancers, such as acute myelogenous leukemia (AML), myelodysplastic syndromes (MDS), and certain other types of cancers of the skin and other organs. Talk with your healthcare provider about your risk of developing new cancers if you take REVLIMID. Your healthcare provider will check you for new cancers during your treatment with REVLIMID.

Please see Important Safety Information throughout and full [Prescribing Information](#), including Boxed WARNINGS and [Medication Guide](#).



Important Safety Information (continued)

What are the possible side effects of REVLIMID® (lenalidomide)? (continued)

- **Severe liver problems, including liver failure and death.** Your healthcare provider should do blood tests to check your liver function during your treatment with REVLIMID. Tell your healthcare provider right away if you develop any of the following symptoms of liver problems:
 - yellowing of your skin or the white part of your eyes (jaundice)
 - dark or brown (tea-colored) urine
 - pain on the upper right side of your stomach area (abdomen)
 - bleeding or bruising more easily than normal
 - feeling very tired
- **Severe skin reactions and severe allergic reactions** can happen with REVLIMID and may cause death.

Call your healthcare provider right away if you develop any of the following signs or symptoms during treatment with REVLIMID:

- a red, itchy skin rash
- peeling of your skin or blisters
- severe itching
- fever

Get emergency medical help right away if you develop any of the following signs or symptoms during treatment with REVLIMID:

- swelling of your lips, mouth, tongue, or throat
- trouble breathing or swallowing
- raised red areas on your skin (hives)
- a very fast heartbeat
- you feel dizzy or faint
- **Tumor lysis syndrome (TLS).** TLS is caused by the fast breakdown of cancer cells. TLS can cause kidney failure and the need for dialysis treatment, abnormal heart rhythm, seizure, and sometimes death. Your healthcare provider may do blood tests to check you for TLS.
- **Worsening of your tumor (tumor flare reaction).** Tell your healthcare provider if you get any of these symptoms of tumor flare reaction while taking REVLIMID: tender, swollen lymph nodes; low-grade fever, pain, or rash.

Please see Important Safety Information throughout and full [Prescribing Information](#), including Boxed WARNINGS and [Medication Guide](#).



Important Safety Information (continued)

What are the possible side effects of REVLIMID® (lenalidomide)? (continued)

- **Thyroid problems.** Your healthcare provider may check your thyroid function before you start taking REVLIMID and during treatment with REVLIMID.
- **Risk of early death in MCL.** In people who have mantle cell lymphoma (MCL), there may be a risk of dying sooner (early death) when taking REVLIMID. Talk with your healthcare provider about any concerns and possible risk factors.

The most common side effects of REVLIMID include:

- diarrhea
- rash
- nausea
- constipation
- tiredness or weakness
- fever
- itching
- swelling of your arms, hands, legs, feet, and skin
- sleep problems (insomnia)
- headache
- muscle cramps or spasms
- shortness of breath
- cough, sore throat, and other symptoms of a cold
- upper respiratory tract infection or bronchitis
- inflammation of the stomach and intestine (“stomach flu”)
- nose bleed
- shaking or trembling (tremor)
- joint aches
- pain in your back or stomach area (abdomen)

These are not all of the possible side effects of REVLIMID. Your healthcare provider may tell you to decrease your dose, temporarily stop or permanently stop taking REVLIMID if you develop certain serious side effects during treatment with REVLIMID. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088.

Please see Important Safety Information throughout and full [Prescribing Information](#), including Boxed WARNINGS and [Medication Guide](#).



Notes:

Use this section to write down any side effects you experience or questions you want to remember to ask your doctor.

Please see Important Safety Information throughout and full [Prescribing Information](#), including [Boxed WARNINGS](#) and [Medication Guide](#).





Join *My Moments, My Support*

Sign up at MyMomentsMySupport.com to receive ongoing information and learn more about your disease and treatment.

Please see Important Safety Information throughout and full [Prescribing Information](#), including Boxed WARNINGS and [Medication Guide](#).





Understanding Side Effects

To learn more about the possible side effects of taking REVLIMID® (lenalidomide), visit [REVLIMID.com](https://www.revlimid.com).

Please see REVLIMID Important Safety Information throughout and on pages 2-15, as well as full [Prescribing Information](#), including **Boxed WARNINGS** and [Medication Guide](#).



REVLIMID® is a registered trademark of Celgene Corporation, a Bristol-Myers Squibb Company.